arXiv:1110.1611v1 [q-bio.PE] 7 Oct 2011

A hierarchical network approach for modeling Rift Valley fever epidemics

Ling Xue[a,*], H. Morgan Scott[b], Lee W. Cohnstaedt[c], Caterina Scoglio[a]

[a]*Department of Electrical & Computer Engineering,*
Kansas State University, U.S. 66506
[b]*Department of Diagnostic Medicine/Pathobiology,*
Kansas State University, U.S. 66506
[c] *Center for Grain and Animal Health Research,*
United States Department of Agriculture, U.S. 66502

Abstract

Rift Valley fever is a vector-borne disease which causes high morbidity and mortality in livestock and also can infect humans. In the event Rift Valley fever virus is introduced to the United States, mitigation strategies need to be rapidly set in place to prevent the disease from becoming endemic. Presented here is a stochastic network-based mathematical model of Rift Valley fever. Given a lack of empirical data for this region, this difference equation model uses stochastic parameters following several Pert distributions. A large directed, asymmetric network was established with 3621 nodes in the Uvalde and Panhandle areas of Texas, U.S.A. noting that the climate in Texas can be similar to that of the Rift Valley in Eastern Africa. This model predicts how Rift Valley fever can spread spatially based on biotic and antibiotic factors and identifies the relevant parameters in RVFV propagation. Humans, cattle, *Aedes* mosquitoes, *Culex* mosquitoes are independently treated with different contact networks to assess virus propagation and spread under different scenarios. The importance of the factors affecting Rift Valley fever virus spread was assessed under different scenarios. Spatial effects and climate factors are also addressed in the model. It has been shown that cattle movement, mosquito population and favorable climate for mosquito development contribute to the spatial spread of Rift Valley fever virus.

Keywords:
Network, Meta-population, Stochastic parameters, Rift Valley fever (RVF), control, mitigation, *Aedes* mosquitoes, *Culex* mosquitoes

---

1. Introduction

Rift Valley fever (RVF) was first identified in Egypt in 1931 [6] and is endemic in east and South Africa [9]. It is a mosquito-borne viral disease which mainly affects ruminants and humans [17]. Rift Valley fever causes abortion in adult and death in newborn livestock [17]. Sheep, goats

---

[*] Corresponding Author

Email addresses: `lxue@ksu.edu` (Ling Xue), `hmscott@vet.k-state.edu` (H. Morgan Scott), `Lee.Cohnstaedt@ars.usda.gov` (Lee W. Cohnstaedt), `caterina@k-state.edu` (Caterina Scoglio)

and cattle are the most important affected animal hosts from an economic standpoint [9]. Humans and domestic livestock are the main hosts [5]. The virus is maintained by vertical transmission within *Aedes* mosquitoes [16]. High levels RVF activity is usually related to persistent, above normal rainfall and ENSO event in East Africa [15].

Ruminants usually acquire the infection from the bites of mosquitoes [18]. *Aedes* and *Culex* mosquitoes are thought of as the main vectors with respect to vector competence [5]. *Aedes* mosquitoes lay eggs in dry mud [30] and the eggs can survive for a long period [9]. After flooding, the eggs infected by Rift Valley fever virus (RVFV) can develop into infected adult mosquitoes [9].

Movement of infected animals is thought of as a possible pathway for introducing RVFV to a disease-free area [19]. In Egypt in 1977 [25], the trade of sheep from east Africa to Mecca in Saudi Arabia or in Yemen during a festival period was considered to be a likely pathway for the introduction of RVFV [2, 7]. As a consequence, a ban on livestock trade can be an effective way to prevent further spread of RVFV [7].

Humans can acquire the infection from the bites of mosquitoes or bodily fluids of animals during butchering or slaughtering [1]. However, most human infections result from contact with animals [9]. Individuals working with animals, such as farmers and veterinarians, are the most vulnerable to RVFV infection [23]. Rift Valley fever virus infection causes severe influenza-like disease in humans with serious consequences such as blindness, or even death [17]. It has been reported that more than 200 humans died of RVF in Mauritania in 1987 [13]. There were 738 reported human cases in Sudan including 230 deaths, in 2007 – 2008 [28]. It is possible that the number of human cases has been underreported, especially in rural areas [5]. Rift Valley fever virus has already spread to Yemen and Saudi Arabia [5] and the French island of Mayotte with multiple human cases reported [26]. Rift Valley fever virus may be introduced to the United States, similar to West Nile virus [10], and the consequences of RVFV transmission to humans in new locations should be focused on.

Rift Valley fever models have been proposed; however, no model addresses spatial transmission during an outbreak [19]. Very few mathematical dynamic transmission models have explored mechanisms of RVFV circulation [19], although one dealt with the varied prevalence of RVF and mosquito abundance data from 1991 to 1996 [4]. A theoretical model in a closed system including *Aedes* and *Culex* mosquitoes and livestock population has been proposed by Gaff et al. [11]. The authors suggest that RVFV can exist if the contact rate between vectors and hosts is high based on sensitivity analysis of the theoretical model parameters [19]. However, neither human hosts nor spatial effects are considered in their model. A theoretical mathematical model on RVF with human hosts has been proposed [20]. The important parameters to the transmission of RVFV have been assessed in their model using sensitivity analysis. Human hosts are considered in their model, but spatial effects and climate factors are not considered. For the Ferlo area of Senegal, a pond-level meta-population model was built assuming that *Aedes* mosquitoes are the only vector and rainfall is the only driving force [9]. It has been shown that within-Ferlo virus is possible to persist if the livestock move between ponds and the rainfall does not occur in all ponds simultaneously [9]. Although various transmission scenarios have been explored, no hosts were considered in the model. No hosts are considered in the model. A deterministic meta-population model including human hosts has been proposed by our group [29] and a study was carried out for South Africa. The model is based on RVFV spatial transmission during an outbreak, where a small network with three nodes has been established [29]. In reality, the propagation of the epidemic is very complicated, so a deterministic model is not always enough to reflect the dynamic spread of the disease. The real network can be large, complicated and hierarchical.

Proposed here is a stochastic network-based mathematical model with four competent species, that is, *Aedes* mosquitoes, *Culex* mosquitoes, cattle and humans. The incidence of infection can be predicted and mitigation strategies can be assessed with this model. Texas is roughly climatically similar to areas of the Rift Valley in eastern Africa where RVF is endemic . It's hypothesized that RVFV can be introduced to Texas based on empirical animal and movement data available for two regions in Texas. The nodes were chosen as 3526 farms in nine counties in South Texas: Bandera, Edwards, Frio, Kinney, Maverick, Medina, Real, Uvalde, and Zavala Counties [27], three livestock markets located in three counties: Uvalde, Medina and Frio counties [27], and 92 feedlots located in the northern Panhandle area [27]. In all, 3621 nodes populated the large network. Different scenarios were then simulated to analyze how RVFV will spread and which factors that are playing an important role in RVFV spreading if infected mosquitoes arrive in the United States. The results have shown that cattle movement, mosquito population, and climate play an important role in the spatial spread of RVFV.

2. The stochastic mathematical model

Three different sets of nodes were considered in the model, namely, 3526 cattle farms, or ranches and 3 livestock markets in the southwest of Texas and 92 feedlots in the northwest part of the state [27]. The nodes were considered homogeneous within each set. This meta-population model is used to study RVF dynamics in the 3621 locations. Each farm, market or feedlot is represented as location i, $i = 1, \cdots, 3621$. All the transitions to be discussed below are for location i at day t.

2.1. *The Difference equation model*

The model is based on the following assumptions.

- Each farm, livestock market, feedlot is a node. The network is directed and asymmetric considering movement in different directions and rates of moving in and out of a node. Farms, feedlots and markets are in different categories respectively.
- The animals include only cattle.
- Once an animal leaves a node, it never returns to that node.
- All animals are homogeneously mixed within a node.
- Only RVFV transmission in farms and in feedlots is considered because markets are hosts-transit nodes.
- The movement rate of cattle in infected compartment is assumed to be half of that of the cattle in other compartment in the absence of further information. Any realistic movement rate can be adopted in the model.
- The mosquitoes will only diffuse to adjacent nodes in a time step. The destination of cattle movement is randomly selected and independent of distance.
- The birth rate and death rate is 0 in markets, birth rate is 0 in feedlots.
- Human birth, death, mortality and mobility are ignored.

- * Birth, death, infection, transfer between compartments, mosquitoes diffusion, and cattle movement occur at the end of each time step.

2.1.1. *Aedes* mosquito vectors

Below is the set of diffence equations for *Aedes* mosquitoes. The parameters and variables are described in 2.2.

$$\begin{aligned}
P_{1i}(t+1) - P_{1i}(t) &= (b_1\left[N_{1i}(t) - q_1 I_{1i}(t)\right] - \theta_1 P_{1i}(t)) * \delta_A(t) && (1)\\
Q_{1i}(t+1) - Q_{1i}(t) &= (b_1 q_1 I_{1i}(t) - \theta_1 Q_{1i}(t) * \delta_A(t) && (2)\\
S_{1i}(t+1) - S_{1i}(t) &= \theta_1 P_{1i}(t) + \sum_{j=1,j\neq i}^{n} \omega_{ji}^{1} S_{1j}(t) - \sum_{j=1,j\neq i}^{n} \omega_{ij}^{1} S_{1i}(t) - d_1 S_{1i}(t) &&\\
&\quad - \beta_{21} S_{1i}(t) I_{2i}(t)/N_{2i}(t) && (3)\\
E_{1i}(t+1) - E_{1i}(t) &= \sum_{j=1,j\neq i}^{n} \omega_{ji}^{1} E_{1j}(t) - \sum_{j=1,j\neq i}^{n} \omega_{ij}^{1} E_{1i}(t) - d_1 E_{1i}(t) &&\\
&\quad + \beta_{21}(t) S_{1i}(t) I_{2i}(t)/N_{2i}(t) - \varepsilon_1 E_{1i}(t) && (4)\\
I_{1i}(t+1) - I_{1i}(t) &= \sum_{j=1,j\neq i}^{n} \omega_{ji}^{1} I_{1j}(t) - \sum_{j=1,j\neq i}^{n} \omega_{ij}^{1} I_{1i}(t) + \theta_1 Q_{1i}(t) - d_1 I_{1i}(t) + \varepsilon_1 E_{1i}(t) && (5)\\
N_{1i}(t+1) &= S_{1i}(t+1) + E_{1i}(t+1) + I_{1i}(t+1) && (6)\\
Q_{1i}(t+1) &= Q_{1i}(T_Q\left\lfloor t/T_Q \right\rfloor + 1) && (7)
\end{aligned}$$

A modified S-E-I model includes compartments P and Q which are uninfected eggs and infected eggs respectively [29]. There are $b_1 N_{1i}(t)$ eggs laid, including $b_1 q_1 I_{1i}(t)$ infected eggs and $b_1 N_{1i}(t) - b_1 q_1 I_{1i}(t)$ uninfected eggs each day. After the development period, $\theta_1 P_{1i}(t)$ uninfected eggs develop into susceptible adult *Aedes* mosquitoes and $\theta_1 Q_{1i}(t)$ infected eggs develop into infected adult *Aedes* mosquitoes every T_Q days. The percentage of *Aedes* mosquitoes moving from location i to location j is ω_{ij}^{1}. The change in the number of *Aedes* mosquitoes due to mobility in compartment X is given as $\sum_{j=1,j\neq i}^{n} \omega_{ji}^{1} X_{1j}(t) - \sum_{j=1,j\neq i}^{n} \omega_{ij}^{1} X_{1i}(t)$. The number of infected *Aedes* mosquitoes from contact with livestock is $\beta_{21} S_{1i}(t) I_{2i}(t)/N_{2i}(t)$. Following the incubation period, $\varepsilon_1 E_{1i}(t)$ *Aedes* mosquitoes transfer from exposed state to infected state. *Aedes* mosquitoes dying naturally in compartment X is given as $d_1 X_{1i}(t)$. Dead mosquitoes are removed from the compartments. If the average temperature is above 60 ^{o}F and the average rainfall is above 0 *mm* during the T_Q days, then $\delta_A(t) = 1$, otherwise $\delta_A(t) = 0$. If the average temperature or rainfall during the period does not reach the threshold, new eggs are not produced and the infected eggs will not develop into adult mosquitoes.

2.1.2. *Culex* mosquito vectors

$$\begin{aligned}
P_{3i}(t+1) - P_{3i}(t) &= b_3(Temp, precipitation, T_d, t) N_{3i}(t) - \theta_3(Temp, t) P_{3i}(t) && (8)\\
S_{3i}(t+1) - S_{3i}(t) &= \theta_3(temp, t) P_{3i}(t) + \sum_{j=1,j\neq i}^{n} \omega_{ji}^{3} S_{3j}(t) - \sum_{j=1,j\neq i}^{n} \omega_{ij}^{3} S_{3i}(t) - d_3 S_{3i}(t) &&\\
&\quad - \beta_{23} S_{3i}(t) I_{2i}(t)/N_{2i}(t) && (9)
\end{aligned}$$

$$\begin{aligned} E_{3i}(t+1) - E_{3i}(t) = &\sum_{j=1,j\neq i}^{n} \omega_{ji}^{3} E_{3j}(t) - \sum_{j=1,j\neq i}^{n} \omega_{ij}^{3} E_{3i}(t) - \varepsilon_3 E_{3i}(t) - d_3 E_{3i}(t) \\ &+ \beta_{23} S_{3i}(t) I_{2i}(t)/N_{2i}(t) \end{aligned} \tag{10}$$

$$I_{3i}(t+1) - I_{3i}(t) = \sum_{j=1,j\neq i}^{n} \omega_{ji}^{3} I_{3j}(t) - \sum_{j=1,j\neq i}^{n} \omega_{ij}^{3} I_{3i}(t) + \varepsilon_3 E_{3i}(t) - d_3 I_{3i}(t) \tag{11}$$

$$N_{3i}(t+1) = S_{3i}(t+1) + E_{3i}(t+1) + I_{3i}(t+1) \tag{12}$$

Above is a modified S-E-I model for *Culex* mosquitoes. Only uninfected eggs are included since the female *Culex* mosquito has not been demonstrated to transmit RVFV to its own eggs [11]. There are $b_3 N_{3i}(t)$ eggs laid each day. After the development period, $\theta_3 P_{3i}(t)$ eggs develop into susceptible adult *Culex* mosquitoes. The percentage of *Culex* mosquitoes moving from location i to location j is ω_{ij}^{3}. The change in the number of *Culex* mosquitoes due to movement in compartment X is given as $\sum_{j=1,j\neq i}^{n} \omega_{ji}^{3} X_{3j}(t) - \sum_{j=1,j\neq i}^{n} \omega_{ij}^{3} X_{3i}(t)$. The number of *Culex* mosquitoes acquiring infection from livestock is denoted by $\beta_{23} S_{3i}(t) I_{2i}(t)/N_{2i}(t)$. After the incubation period, $\varepsilon_3 E_{3i}(t)$ *Culex* mosquitoes transfer to the infected compartment. The *Culex* mosquitoes removed from the compartment X due to natural death is $d_3 X_{3i}(t)$. The total number of *Culex* mosquitoes is also variable and equals the sum of the *Culex* mosquitoes in all the compartments.

2.1.3. Livestock hosts

$$\begin{aligned} S_{2i}(t+1) - S_{2i}(t) = &\, b_2(t)\delta_b(i) N_{2i}(t) + \sum_{j=1,j\neq i}^{n} \omega_{ji}^{2} S_{2j}(t) - \sum_{j=1,j\neq i}^{n} \omega_{ij}^{2} S_{2i}(t) - d_2 \delta_d(i) S_{2i}(t) N_{2i}(t)/K_2 \\ &- \beta_{12} S_{2i}(t) I_{1i}(t)/N_{1i}(t) - \beta_{32} S_{2i}(t) I_{3i}(t)/N_{3i}(t) \end{aligned} \tag{13}$$

$$\begin{aligned} E_{2i}(t+1) - E_{2i}(t) = &\sum_{j=1,j\neq i}^{n} \omega_{ji}^{2} E_{2j}(t) - \sum_{j=1,j\neq i}^{n} \omega_{ij}^{2} E_{2i}(t) - d_2 \delta_d(i) E_{2i}(t) N_{2i}(t)/K_2 - \varepsilon_2 E_{2i}(t) \\ &+ \beta_{12} S_{2i}(t) I_{1i}(t)/N_{1i}(t) + \beta_{32} S_{2i}(t) I_{3i}(t)/N_{3i}(t) \end{aligned} \tag{14}$$

$$\begin{aligned} I_{2i}(t+1) - I_{2i}(t) = &\sum_{j=1,j\neq i}^{n} \omega_{ji}^{2} I_{2j}(t) - \sum_{j=1,j\neq i}^{n} \omega_{ij}^{2} I_{2i}(t) - d_2 \delta_d(i) I_{2i}(t) N_{2i}(t)/K_2 + \varepsilon_2 E_{2i}(t) \\ &- \gamma_2 I_{2i}(t) - \mu_2 I_{2i}(t) \end{aligned} \tag{15}$$

$$R_{2i}(t+1) - R_{2i}(t) = \sum_{j=1,j\neq i}^{n} \omega_{ji}^{2} R_{2j}(t) - \sum_{j=1,j\neq i}^{n} \omega_{ij}^{2} R_{2i}(t) + \gamma_2 I_{2i}(t) - d_2 \delta_d(i) R_{2i}(t) N_{2i}(t)/K_2 \tag{16}$$

$$N_{2i}(t+1) = S_{2i}(t+1) + E_{2i}(t+1) + I_{2i}(t+1) + R_{2i}(t+1) \tag{17}$$

It's assumed that cattle can be transported from farms to markets or feedlots, or from markets to farms or feedlots, but not from feedlots to farms. The total number of newborn livestock in location i is $b_2(t) N_{2i}(t)$ each day. The percentage of cattle moving from location i to location j is ω_{ij}^{2}. The change in the number of livestock in compartment X due to mobility is given as $\sum_{j=1,j\neq i}^{n} \omega_{ji}^{2} X_{2j}^{[m]}(t) - \sum_{j=1,j\neq i}^{n} \omega_{ij}^{2} X_{2i}^{[m]}(t)$. After the incubation period, $\varepsilon_2 E_{2i}(t)$ livestock transfer from exposed state to infected state. The number of cattle infected by *Aedes* mosquitoes and *Culex* mosquitoes is denoted as $\beta_{12} S_{2i}(t) I_{1i}(t)/N_{1i}(t)$ and $\beta_{32} S_{2i}(t) I_{3i}(t)/N_{3i}(t)$, respectively. After the infectious period, $\gamma_2 I_{2i}(t)$ cattle recover from RVFV infection. The number of dead cattle in each compartment X is given as $d_2 X_{2i} N_{2i}(t)/K_2$ in which K_2 is the carrying capacity of cattle in each node. There are $\mu_2 I_{2i}(t)$ cattle dying of RVFV infection. The variables $\delta_b(i)$ and $\delta_d(i)$ are used to differentiate different type of nodes. If location i is a farm, then $\delta_b(i) = 1$, $\delta_d(i) = 1$. If location i is a market, then $\delta_b(i) = 0$, $\delta_d(i) = 0$. If location i is a feedlot, then $\delta_b(i) = 0$.

2.1.4. Human hosts

$$S_{4i}(t+1) - S_{4i}(t) = -\beta_{14} S_{4i}(t) I_{1i}(t)/N_{1i}(t) - \beta_{24} S_{4i}(t) I_{2i}(t)/N_{2i}(t) - \beta_{34} S_{4i}(t) I_{3i}(t)/N_{3i}(t) \tag{18}$$

$$\begin{aligned} E_{4i}(t+1) - E_{4i}(t) = {} & \beta_{14} S_{4i}(t) I_{1i}(t)/N_{1i}(t) + \beta_{24} f_i S_{4i}(t) I_{2i}(t)/N_{2i}(t) + \beta_{34} S_{4i}(t) I_{3i}(t)/N_{3i}(t) \\ & - \varepsilon_4 E_{4i}(t) \end{aligned} \tag{19}$$

$$I_{4i}(t+1) - I_{4i}(t) = \varepsilon_4 E_{4i}(t) - \gamma_4 I_{4i}(t) \tag{20}$$

$$R_{4i}(t+1) - R_{4i}(t) = \gamma_4 I_{4i}(t) \tag{21}$$

Assume that the susceptible human population only includes those working with animals, such as veterinarians and farmers, because they are at higher risk of RVFV exposure and infection [23]. The number of humans infected by *Aedes* mosquitoes, *Culex* mosquitoes, and livestock is $\beta_{14} S_{4i}(t) I_{1i}(t)/N_{1i}(t)$, $\beta_{24} S_{4i}(t) I_{2j}(t)/N_{2i}(t)$, and $\beta_{34} S_{4i}(t) I_{3i}(t)/N_{3i}(t)$ respectively. There are $\varepsilon_4 E_{4i}(t)$ humans transferring to the infected compartment after the incubation period and $\gamma_4 I_{4i}(t)$ humans recover from RVFV infection.

2.2. Descriptions of parameters and variables in the model

The descriptions of the parameters of the following equations are listed in Table 1.
The variables in the equations are as following:
$P_{1i}(t)$=The number of uninfected *Aedes* mosquito eggs in location i
$Q_{1i}(t)$=The number of infected *Aedes* mosquito eggs in location i
$S_{1i}(t)$=The number of susceptible *Aedes* mosquitoes in location i
$E_{1i}(t)$=The number of exposed *Aedes* mosquitoes in location i
$I_{1i}(t)$=The number of infected *Aedes* mosquitoes in location i
$N_{1i}(t)$=The total number of *Aedes* mosquitoes in location i
$S_{2i}(t)$=The number of susceptible livestock in location i
$E_{2i}(t)$=The number of exposed livestock in location i
$I_{2i}(t)$=The number of infected livestock in location i
$R_{2i}(t)$=The number of recovered livestock in location i
$N_{2i}(t)$=The total number of livestock in location i
$P_{3i}(t)$=The number of uninfected *Culex* mosquito eggs in location i
$S_{3i}(t)$=The number of susceptible *Culex* mosquitoes in location i
$E_{3i}(t)$=The number of exposed *Culex* mosquitoes in location i
$I_{3i}(t)$=The number of infected *Culex* mosquitoes in location i
$N_{3i}(t)$=The total number of *Culex* in location i
$S_{4i}(t)$=The number of susceptible humans in location i
$E_{4i}(t)$=The number of exposed humans in location i
$I_{4i}(t)$=The number of infected humans in location i
$R_{4i}(t)$=The number of recovered humans in location i

3. The network

The movement between the three sets of nodes is shown in Figure 2(a). The network is directed and asymmetric. The distance between any two farms is compared with an assumed radius which is defined as 2 *km* here. If the distance is smaller than the radius, then the two farms are neighbors for mosquito movement and there is a link between them. An example of the mosquito movement network is shown in Figure 2(b). All nodes are shown in Figure 3(a), where links are omitted due to their large number. An example of the links of a single farm is shown in Figure 3(b). The exact location of each farm, market and feedlot is obscured because those data are not available to public. The nodes are uniformly distributed in each county according to the total number of farms in each county [27]. Some markets are randomly assigned as neighbors of each farm and some farms are randomly assigned as neighbors of each market. The neighbors are the destination of the cattle. Farms and markets are thought of as sources of cattle to feedlots. The destination, markets or feedlots, from cattle farm is independent of distance as are cattle from markets. The movement rate is inverse to distance. The nodes are distributed in each county according to the total number of feedlots in each county in the Panhandle area [27]. The initial number of susceptible cattle in each farm or feedlot for numerical simulation is assumed according to the total number of cattle in each county [27]. The color of the node is determined by the number of infected cattle in each farm, market or feedlot. Green represents the susceptible cattle and humans respectively. Initially, there are no infected nodes in any of the counties.

4. Case Study: Texas, U.S.A. 2011

Case study was carried out for Texas, U.S.A from January, 2005 to December, 2006. The parameters of the above equations are listed in Table 1. The parameters with given ranges are variables following severals Pert distributions. Uniform distribution has equal probability for any value within the range. Pert distribution has a smooth trend and flat tail as Gaussian distribution as well as a flexible most possible value. Therefore, Pert distribution is chosen for our preference. The movement rates of livestock are shown in Table 2. The movement rate is assumed to be the maximum percentage of animals moving [8]. The number of farms and cattle in each of the nine counties [21] is shown in Table 3.

where:
$n_m(i)$=the number of markets connected to farm i
$n_f(i)$=the number of farms connected to market i
$n_{ffe}(i)$=the number of feedlots connected to farm i
$n_{mfe}(i)$=the number of feedlots connected to market i
dis_{ij}=the distance between node i and node j

The diffusion rates of *Aedes* and *Culex* mosquitoes are shown below [24].

$$\omega_{ij}^1 = \omega_{ij}^3 = \begin{cases} 0 & \text{if the nodes are disjoint} \\ diff/d_{ij}^2 & \text{if two nodes share a border} \end{cases}$$

where $d_{ij}$is the distance between the centers of node i and node j [24] and $diff$ is a diffusion like parameter within the range $(830, 8300)m^2/\text{day}$ [24].

The development rate and egg laying rate of *Culex* mosquitoes [12] are shown in Equation (22) and Equation (23). The parameters adopted are derived from data concerning West Nile



(a) Egg laying rate of *Culex* mosquitoes (b) Birth rate of *Culex* mosquitoes



(c) Birth rate of *Culex* mosquitoes in Texas from January, 2005 to December, 2006 (d) Development rate of *Culex* mosquitoes in Texas from January, 2005 to December, 2006

Figure 1: The relationship between b_3, θ_3 and climate



(a) Animal movement location (b) An example of mosquito network

Figure 2: Movement diagrams



(a) All nodes in Panhandle and Uvalde area (b) The links of a farm

Figure 3: Network in Panhandle and Uvalde area of Texas

virus in 2010 in the northern US [24]. This is the best model currently available for us at the moment. Any more realistic models or parameters can later be adopted in the model once available. The egg laying rate changes with moisture [12] and development rate changes with temperature [12] are shown in Figure 1(a) and Figure 1(b) , respectively. The egg laying rate and development rate of *Culex* mosquitoes in Texas from January, 2005 to December, 2006 are shown in Figure 1(c) and Figure 1(d), respectively. The development rate in Figure 1(d) is enlarged 100 times to be seen clearly. As is shown, the development rate of the *Culex* mosquitoes changes with temperature. The colder it is, the fewer *Culex* eggs are produced. If the temperature is too low, the eggs won't develop into adult mosquitoes. If the temperature is too high, the lifespan of the mosquitoes is shortened so that the development rate decreases. Moisture index is the difference between precipitation and evaporation as shown in Equation (24). The lower the moisture index, the fewer adult mosquitoes are produced because low moisture index represents low precipitation and high evaporation. For some days, the missing precipitation data from January, 2005 to December, 2006 [22] missing are assumed to be zero. The evaporation data are calculated using the formula in Equation (25) [14]. One set of climate data was applied to the Uvalde study area and another set of climate data was applied to Panhandle area. The rainfall and temperature determine the presence of the mosquitoes. They are necessary for mosquitoes to survive and breed. The presence of the infected eggs is clearly important to begin the simulated epidemics.

$$\theta_3(Temp,t) = A * \frac{(Temp(t)+K)}{298.15} * \frac{exp[\frac{HA}{1.987} * (\frac{1}{298.15} - \frac{1}{Temp(t)+K})]}{1 + exp[\frac{HH}{1.987} * (\frac{1}{TH} - \frac{1}{Temp(t)+K})]} \tag{22}$$

$$b_3(Temp, precipitation, T_d, t) = b_0 + \frac{Emax}{1 + exp[-\frac{Moisture(t)-Emean}{Evar}]} \tag{23}$$

$$Moisture(t) = \sum_{D=t-6}^{t} precipitation(D) - Evaporation(D) \tag{24}$$

$$\begin{aligned} Evaporation(t) &= \frac{700(Temp(t)+0.006h)/(100-latitude)}{80-Temp(t)} \\ &+ \frac{15(Temp(t)-T_d(t))}{80-Temp(t)} mm/day \end{aligned} \tag{25}$$

Where:

$Temp(t)$ =air temperature in units of ^oC [14]
$latitude$ =the latitude (degrees) [14]
$T_d(t)$ =the mean dew-point in units of ^oC [14]
h =the elevation (meters) [14]

Various scenarios have been examined and listed in Table 4. Simulations were run with 200 different sets of parameters using a daily time step. The climate data from January, 2005 to December, 2006 [22] in Texas was adapted into the model. The spread of RVFV with time is shown in Figure 4. The parameter T_Q is set to 15 as an example. Digital numbers are used to distinguish farms. As an illustration, the distance between Farm 3526 and other 4 farms is less than 2 *km*. So Farm 3526 has 4 neighbors. The change in the number of infected cattle or humans in farms during an outbreak is the main focus of this case study. Therefore, the number of infected cattle and humans in feedlots are not illustrated in the following figures. Assume 10 infected *Aedes* mosquito eggs are introduced to Farm 3526. The number of infected humans and cattle in Farm 3526 and its neighbor farms is shown in Figure 4(a) and Figure 4(b). To see the number of infected cattle and humans more clearly, the infected humans and animals only in the neighbor farms of Farm 3526 is shown in Figure 4(c) and Figure 4(d). If there are no infected eggs in all nodes at the beginning, then the percentage of infected animals and humans is zero in all farms. Farm 3440 is one of the neighbors of Farm 3526. The percentage of infected cattle and humans in the neighbor nodes of Farm 3440 shown in Figure 4(e) and Figure 4(f). The infected cattle and humans also arise in the neighbors of Farm 3440. If there is no movement, then there are no infected cattle or humans in the neighbors of Farm 3526 as shown in Figure 4(g) and Figure 4(h). The spatial spread of RVFV with 10000 and 10 eggs in a small farm of Bandera county initially is shown in Figure 5. The percentage of the infected cattle and humans reaches the peak in March for Farm 3526 and around May for its neighbors. Rift Valley fever virus spreads much more quickly with more infected eggs initially as shown in the figure. If 10 infected eggs only exist in a farm with very few cattle or humans at the beginning, then cases likely won't spread quickly to other counties as shown in Figure 5(a) and Figure 5(b). Therefore, the infection of a small farm with few infected eggs does not affect other nodes like an epidemic in a major node. The spatial spread of RVFV with 10 infected eggs in a large farm of Zavala county initially is shown in Figure 6. The percentage of infected cattle and humans on the $100th$ day is shown in Figure 6(a) and Figure 6(b) respectively. By way of more comparison of Figure 5 and Figure 6, Rift Valley fever virus spreads more quickly with infected eggs in a larger farm initially. The percentage of infected cattle and humans on the $100th$ day with infected eggs in 9 large farms of 9 counties at the beginning is shown in Figure 7(a) and Figure 7(b). RVFV has spread to other counties much more quickly as shown in the figures because more counties have infected eggs at the beginning. The infected cattle and humans also arise in the nodes of other counties because of livestock movement. If infected eggs are in large farms of different counties, RVFV will spread quickly. If 10 infected eggs are only in Farm 3526 at the beginning and there is only mosquito movement, the percentage of infected cattle and humans on the $100th$ day is shown in Figure 8. If 10 infected eggs are only in Farm 3526 at the beginning and there is only cattle movement, the percentage of infected cattle and humans on the $100th$ day is shown in Figure 9. The spatial spread of RVFV is more rapid with cattle movement than with only mosquito movement. The percentage of infected cattle and humans being infected with mosquito movement only, and 10 and 10000 infected eggs initially, are shown in Figures 10(a), 10(b), 10(c), 10(d) and Figure 10(e), 10(f), 10(g), 10(h), respectively. As is shown in Figure 10, the number of infected eggs greatly affects the percentage of infected cattle and humans. The percentage of infected cattle and humans changing with time with only cattle movement and with different numbers of infected eggs are shown in Figure 11. Therefore, the movement of cattle contributes to the spatial spread of RVFV with the given initial condition and movement rate as shown in Figure 8, Figure 9, Figure 10 and Figure 11. This gives us some information on potential for mitigation of RVFV spread. Banning animal trade and movement may help prevent further spread of RVFV during an outbreak.

5. Discussions

Presented here is a mathematical model with stochastic parameters describing the spatial spread of RVFV during an outbreak. The network is a large hierarchical network with three different sets of nodes which are farms, markets and feedlots. The nodes are homogeneous within

Parameter	Description	Range	Most possible value	Units	Source
β_{12}	adequate contact rate: *Aedes* to livestock	$(0.0021, 0.2762)$	0.1392	1/day	[11]
β_{21}	adequate contact rate: livestock to *Aedes*	$(0.0021, 0.2429)$	0.1225	1/day	[11]
β_{23}	adequate contact rate: livestock to *Culex*	$(0.0000, 0.3200)$	0.16	1/day	[11]
β_{32}	adequate contact rate: *Culex* to livestock	$(0.0000, 0.096)$	0.04	1/day	[11]
β_{14}	adequate contact rate: *Aedes* to humans	$(0.001, 0.002)$	0.0015	1/day	Assume
β_{24}	adequate contact rate: livestock to humans	$(0.00004, 0.00008)$	0.0006	1/day	Assume
β_{34}	adequate contact rate: *Culex* to humans	$(0.0005, 0.001)$	0.00525	1/day	Assume
$1/\gamma_2$	recover rate in livestock	$(2, 5)$	3.5	1/day	[11]
$1/\gamma_4$	recover period in humans	$(4, 7)$	5.5	1/day	[3]
$1/d_1$	longevity of *Aedes* mosquitoes	$(3, 60)$	31.5	days	[11]
$1/d_2$	longevity of livestock	$(360, 3600)$	1980	days	[11]
$1/d_3$	longevity of *Culex* mosquitoes	$(3, 60)$	31.5	days	[11]
b_1	number of *Aedes* eggs laid per day	$1/(3, 60)$	1/31.5	1/day	[11]
b_2	daily birthrate of livestock	$1/(360, 3600)$	1/1980	1/day	[11]
b_3	number of *Culex* eggs laid per day	$1/(3, 60)$	1/31.5	1/day	[11]
$1/\epsilon_1$	incubation period in *Aedes* mosquitoes	$(4, 8)$	6	days	[11]
$1/\epsilon_2$	incubation period in livestock	$(2, 6)$	4	days	[11]
$1/\epsilon_3$	incubation period in *Culex* mosquitoes	$(4, 8)$	6	days	[11]
$1/\epsilon_4$	incubation period in humans	$(2, 6)$	4	days	[20]
μ_2	mortality rate in sheep	$(0.025, 0.1)$	0.0375	1/day	[11]
q_1	transovarial transmission rate in livestock	$(0, 0.1)$	0.05	1/day	[11]
$1/\theta_1$	development time of *Aedes*	$(5, 15)$	10	days	[11]
$1/\theta_3$	development time of *Culex*			days	[11]
K_2	carrying capacity of livestock	100000			Assume
N_f	total number of farms	3526			[21]
N_s	total number of feedlots	92			[21]
N_m	total number of markets	3			[21]
T_Q	development cycle of infected *Aedes* eggs				
A	parameter of the development rate	0.25			[12]
HA	parameter of the development rate	28094			[12]
HH	parameter of the development rate	35692			[12]
TH	parameter of the development rate	298.6			[12]
b_0	minimum constant fecundity rate	0	1/day		[12]
$Emax$	maximum daily egg laying rate	20	1/day		[12]
$Emean$	the mean of the daily egg laying rate	0	1/day		[12]
$Evar$	variance of function	12	1/day		[12]
K	Kelvin parameter	273.15	Kelvin		[12]

Table 1: Parameter ranges for numerical simulation

i	j	range	source
farm	market	60.7% $/(n_m(i) * d_{ij})$	[8]
market	farm	60.7% $/(n_f(i) * d_{ij})$	[8]
farm	feedlot	10.9% $/(n_{ffe}(i) * d_{ij})$	[8]
market	feedlot	10.9% $/(n_{mfe}(i) * d_{ij})$	[8]
feedlot	farm	0	[8]
feedlot	market	0	[8]

Table 2: Movement rate of livestock ω^2_{ij}

County	Cattle and calves	cattle farms	source
Bandera	$11,254$	496	[27]
Edwards	$22,247$	264	[27]
Frio	$51,411$	471	[27]
Kinney	$11,672$	125	[27]
Maverick	$25,135$	183	[27]
Medina	$55,759$	1364	[27]
Real	$4,830$	133	[27]
Uvalde	$52,366$	316	[27]
Zavala	$66,641$	174	[27]
Total		3526	[27]

Table 3: The number of farms and cattle in each county of Uvalde study area in 2007

Scenarios	Effect
No infected *Aedes* eggs in any nodes	No infected cattle or humans in any nodes
Only 10 infected eggs are in a large farm in Zavala county at the beginning	Infected cattle or humans appeared in other counties
Only 10 infected eggs are in farm 1 (a small farm in Bandera county) at the beginning	No infected cattle or humans observed in its neighbors.
The infected eggs are in 9 farms of 9 different counties at the beginning	Infected cattle or humans appeared in other counties
Initially infected eggs are in a large farm in Zavala county and considering mosquito diffusion only	Infected cattle and humans spread to other nodes
Initially infected eggs are in a large farm in Zavala county and considering cattle movement only	Infected cattle and humans spread to other nodes
No cattle movement or mosquito diffusion and infected eggs are in Farm 3526 at the beginning	No spatial spread of infected cattle and humans

Table 4: Scenarios with the climate selected during the most favorable period for mosquito development



(a) Percentage of infected cattle in Farm 3526 and its neighbors

(b) Percentage of infected humans in Farm 3526 and its neighbors



(c) Percentage of infected cattle in the neighbors of Farm 3526

(d) Percentage of infected humans in the neighbors of Farm 3526



(e) Percentage of the infected cattle in the neighbor nodes of one of the neighbors of Farm 3526

(f) Percentage of infected humans in the neighbor nodes of one of the neighbors of Farm 3526



(g) Percentage of infected cattle in Farm 3526 and its neighbors

(h) Percentage of infected humans in Farm 3526 and its neighbors

Figure 4: Numerical simulation results with $T_Q = 15$ where the values are averaged over 200 realizations

each set. The spread of RVFV is estimated not only within farms but also between farms, markets, and feedlots. Evaluating several likely scenarios, the most important factors are identified. Model results indicate

- Local transmission and trans-location transmission are important in accelerating the spread of RVFV.
- The mosquito population and favorable climate for mosquito development contributes to the spread of RVFV among farms.
- The route of moving from farms to livestock markets and subsequently to different farms expands the area of infection.

The existence of infected eggs in large farms contribute to the rapid spread of RVFV. The model is a general model into which parameters in any location can be easily adapted. Climate factors are also involved and analyzed. The diffusion of mosquitoes and the trans-location movement of livestock contribute to the spatial spread of RVFV. Starting with infected eggs in one large node, the infection of RVFV arises in more and more nodes later on. The infected *Aedes* mosquitoes diffuse to adjacent farms and hatch infected offspring. The cattle can be infected by being fed on by *Aedes* mosquitoes. *Culex* mosquitoes amplify the infection following blood meals on infected cattle. The infected cattle transported to markets or feedlots are seeds of the infection in markets or feedlots. Similarly, the virus can be introduced to farms by purchasing infected cattle from markets. The parameters for the development rate and egg laying rate are not uniquely measured for Texas during 2005 to 2006. Rift Valley fever virus can spread quickly to the whole network once it is introduced into a large node. The actual animal movement rate from farms to markets, from markets to farms, from farms to feedlots, from markets to feedlots are not available to us due to confidentiality. Therefore, the incidence data can be inaccurate; however, the model gives us an idea of the pathways for rapid spread of RVFV among farms and counties. It shows that livestock trade and movements play an important role in RVFV spread between counties. restricting livestock trade and the movement of cattle during a RVF outbreak will reduce transmission.



(a) Percentage of infected cattle on the 100*th* day

(b) Percentage of infected humansd on the 100*th* day



(c) Percentage of infected cattle on the 100*th* day

(d) Percentage of infected humans on the 100*th* day

Figure 5: The dynamics of nodes in Uvalde study area with 10 and 10000 infected eggs in a small farm of Bandera county respectively at the beginning of the simulation



(a) Percentage of infected cattle on the 100*th* day

(b) Percentage of infected humans on the 100*th* day

Figure 6: The dynamics of nodes in Uvalde area with 10 infected eggs in a single large farm of Zavala county initially



(a) Percentage of infected cattle on the 100*th* day

(b) Percentage of infected humans on the 100*th* day

Figure 7: The dynamics of nodes in Uvalde area with 10 infected eggs in 9 large farms of 9 counties initially



(a) Percentage of infected cattle on the 100*th* day

(b) Percentage of infected humans the 100*th* day

Figure 8: The dynamics of nodes in Uvalde area with only mosquito diffusion and 10 infected eggs in a large farm of Zavala county initially



(a) Percentage of infected cattle on the 100*th* day

(b) Percentage of infected humans on the 100*th* day

Figure 9: The dynamics of nodes in Uvalde area with only cattle movement and infected eggs in a large farm of Zavala county initially



(a) Percentage of infected cattle in Farm 3526 and its neighbors with 10 infected eggs in Farm 3526 of Zavala county at the beginning considering mosquito diffusion only



(b) Percentage of infected humans in Farm 3526 and its neighbors with 10 infected eggs in Farm 3526 of Zavala county at the beginning considering mosquito diffusion only



(c) Percentage of infected cattle in the neighbors of Farm 3526 with 10 infected eggs in Farm 3526 of Zavala county at the beginning considering mosquito diffusion only



(d) Percentage of infected humans in the neighbors of Farm 3526 with 10 infected eggs in Farm 3526 of Zavala county at the beginning considering mosquito diffusion only



(e) Percentage of infected cattle in Farm 3526 and its neighbors with 10000 infected eggs in a large farm of Zavala county at the beginning considering mosquito diffusion only



(f) Percentage of infected humans in Farm 3526 and its neighbors with 10000 infected eggs in a large farm of Zavala county at the beginning considering mosquito diffusion only



(g) Percentage of infected cattle in the neighbors of Farm 3526 with 10000 infected eggs in Farm 3526 of Zavala county at the beginning considering mosquito diffusion only

(h) Percentage of infected humans in the neighbors of Farm 3526 with 10000 infected eggs in Farm 3526 of Zavala county at the beginning considering mosquito diffusion only

Figure 10: The temporal dynamics of nodes in Uvalde area only with mosquito movement. The values are the aveage over 200 runs.

Acknowledgments

This work has been supported by the DHS Center of Excellence for Emerging and Zoonotic Animal Diseases (CEEZAD), and the National Agricultural Biosecurity Center (NABC) at Kansas State University. We are grateful to Bo Norby, Doyle Fuchs, Bryanna Pockrandt, Livia Olsen, Regina M. Beard, Phillip Schumm, and Annie Gatewood-Hoen for their help in producing this work.

References

[1] European Food Safety Authority (EFSA), . Opinion of the scientific panel on animal health and welfare (ahaw) on a request from the commission related to the risk of a Rift Valley fever incursion and its persistence within the community. `http://www.efsa.europa.eu/fr/efsajournal/doc/238.pdf`.

[2] Abdo-Salem, S., Waret-Szkuta, A., Roger, F., Olive, M.M., Saeed, K., Chevalier, V., 2011. Risk assessment of the introduction of rift Valley fever from the horn of africa to yemen via legal trade of small ruminants. Tropical animal health and production 43, 471–480.

[3] Ahmed, O., . Rift Valley fever A Resurgent Threat Case Studies from Sudan and the Kingdom of Saudi Arabia .

[4] Bicout, D.J., Sabatier, P., 2004. Mapping Rift Valley fever vectors and prevalence using rainfall variations. Vector borne and zoonotic diseases (Larchmont, N.Y.) 4, 33–42.

[5] Chevalier, V., Pepin, M., Plee, L., Lancelot, R., 2010. Rift Valley fever–a threat for europe? Euro surveillance : bulletin europeen sur les maladies transmissibles = European communicable disease bulletin 15, 19506.

[6] Daubney, R., Hudson, J.R., Garnham, P.C., 1931. Enzootic hepatitis or Rift Valley fever. an undescribed virus disease of sheep, cattle and man from east africa. Journal of Pathology and Bacteriology 34, 545–579.

[7] Davies, F.G., 2006. Risk of a Rift Valley fever epidemic at the haj in mecca, saudi arabia. Revue scientifique et technique (International Office of Epizootics) 25, 137–147.

[8] Dominguez, B.J., . Characterization of livestock herds in extensive agricultural settings in southwest texas. `http://repository.tamu.edu/bitstream/handle/1969.1/ETD-TAMU-1302/DOMINGUEZ-THESIS.pdf?sequence=1`.

[9] Favier, C., Chalvet-Monfray, K., Sabatier, P., Lancelot, R., Fontenille, D., Dubois, M., 2006. Rift Valley fever in West Africa: the role of space in endemicity. Tropical Medicine & International Health 11, 1878–1888.

[10] Gaff, H., Burgess, C., Jackson, J., Niu, T., Papelis, Y., Hartley, D., 2011. Mathematical model to assess the relative effectiveness of rift valley fever countermeasures. International Journal of Artificial Life Research (IJALR) 2, 1–18.



(a) Percentage of infected cattle in Farm 3526 and its neighbors with 10 infected eggs in Farm 3526 of Zavala county at the beginning considering cattle movement only



(b) Percentage of infected humans in Farm 3526 and its neighbors with 10 infected eggs in Farm 3526 of Zavala county at the beginning considering cattle movement only



(c) Percentage of infected cattle in the neighbors of Farm 3526 with 10 infected eggs in Farm 3526 of Zavala county at the beginning considering cattle movement only



(d) Percentage of infected humans in the neighbors of Farm 3526 with 10 infected eggs in Farm 3526 of Zavala county at the beginning considering cattle movement only



(e) Percentage of infected cattle in Farm 3526 and its neighbors with 10000 infected eggs in a large farm of Zavala county at the beginning considering cattle movement only



(f) Percentage of infected humans in Farm 3526 and its neighbors with 10000 infected eggs in Farm 3526 of Zavala county at the beginning considering cattle movement only



(g) Percentage of infected cattle in the neighbors of Farm 3526 with 10000 infected eggs in Farm 3526 of Zavala county at the beginning considering cattle movement only

(h) Percentage of infected humans in the neighbors of Farm 3526 with 10000 infected eggs in Farm 3526 of Zavala county at the beginning considering cattle movement only

Figure 11: The temporal dynamics of nodes in Uvalde area with only cattle movement. The values are the aveage over 200 runs.

[11] Gaff, H., Hartley, D., Leahy, N., 2007. An epidemiological model of Rift Valley fever. Electronic Journal of Differential Equations 2007, 1–12.

[12] Gong, H., DeGaetano, A., Harrington, L., 2010. Climate-based models for West Nile Culex mosquito vectors in the Northeastern US. International Journal of Biometeorology , 1–12.

[13] Jouan, A., Guenno, B.L., Digoutte, J.P., Philippe, B., Riou, O., Adam, F., 1988. An RVF epidemic in southern mauritania. Annales de l'Institut Pasteur.Virology 139, 307–308.

[14] Linacre, E., 1977. A simple formula for estimating evaporation rates in various climates, using temperature data alone. Agricultural Meteorology 18, 409–424.

[15] Linthicum, K, J., Anyamba, A., Tucker, C., Kelley, P., Myers, M., C.J., P., 1999. Climate and satellite indicators to forecast rift valley fever epidemics in Kenya. Science 285, 397–400.

[16] Linthicum, K.J., Davies, F.G., Kairo, A., Bailey, C.L., 1985. Rift Valley fever virus (family bunyaviridae, genus phlebovirus). isolations from diptera collected during an inter-epizootic period in Kenya. The Journal of hygiene 95, 197–209.

[17] Martin, V., Chevalier, V., Ceccato, P., Anyamba, A., Simone, L.D., Lubroth, J., de La Rocque, S., Domenech, J., 2008. The impact of climate change on the epidemiology and control of Rift Valley fever. Revue scientifique et technique (International Office of Epizootics) 27, 413–426.

[18] McIntosh, B.M .and Jupp, P., 1981. Epidemiological aspects of rift Valley fever in south africa with reference to vectors. Contrib Epidemiol Biostat 1981, 92–99.

[19] Metras, R., Collins, L.M., White, R.G., Alonso, S., Chevalier, V., Thuranira-McKeever, C., Pfeiffer, D.U., 2011. Rift valley fever epidemiology, surveillance, and control: what have models contributed? Vector borne and zoonotic diseases (Larchmont, N.Y.) 11, 761–771.

[20] Mpeshe, S., Haario, H., Tchuenche, J., . A mathematical model of Rift Valley fever with human host. Acta Biotheoretica , 1–20.

[21] National Agricultural Statistics Service, . 2007 cencus report. `http://www.agcensus.usda.gov/Publications/2007/Full_Report/index.asp`. Accessed March 10, 2010.

[22] National Climatic center, . Noaa satellite and information service. `http://www7.ncdc.noaa.gov/CDO/country`. Accessed Nov 22, 2010.

[23] National Institute for Communicable Diseases, . Interim report on the Rift Valley fever (RVF) outbreak in south africa. `http://www.nicd.ac.za/outbreaks/rvf/docs/RVF_Interim_Report_2010_10_01.pdf`. Accessed Nov 23, 2010.

[24] Otero, M., Solari, H., 2010. Stochastic eco-epidemiological model of dengue disease transmission by Aedes aegypti mosquito. Mathematical biosciences 223, 32–46.

[25] Sellers, R.F., Pedgley, D.E., Tucker, M.R., 1982. Rift Valley fever, egypt 1977: disease spread by windborne insect vectors? The Veterinary record 110, 73–77.

[26] Sissoko, D., Giry, C., Gabrie, P., Tarantola, A., Pettinelli, F., Collet, L., DOrtenzio, E., Renault, P., Pierre, V., 2009.

Rift Valley fever, Mayotte, 2007–2008. Emerging infectious diseases 15, 568.

[27] United States Department of Agriculture, . 2007 census publications. `http://www.agcensus.usda.gov/Publications/2007/Full_Report/Census_by_State/Texas/index.asp`. Accessed March 15, 2011.

[28] World Health Organization, . Rift Valley fever in sudan–update 4. `http://www.who.int/csr/don/2007_11_05/`. Accessed June 23, 2011.

[29] Xue, L., Scott, H.M., Scoglio, C., 2010. A Network-Based Meta-Population Approach to Model Rift Valley Fever Epidemics. ArXiv e-prints `1012.1684`.

[30] Zeller, H., Fontenille, D., Traore-Lamizana, M., Thiongane, Y., Digoutte, J., 1997. Enzootic activity of Rift Valley fever virus in senega. American Journal of Tropical Medicine and Hygiene 56, 265–272.